

# WulffPack: A Python package for Wulff constructions

J. Magnus Rahm<sup>1</sup> and Paul Erhart<sup>1</sup>

<sup>1</sup> Department of Physics, Chalmers University of Technology, Gothenburg, Sweden

DOI: [10.21105/joss.01944](https://doi.org/10.21105/joss.01944)

## Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

---

**Editor:** [Vincent Knight](#) ↗

## Reviewers:

- [@EduPH](#)
- [@mzszym](#)
- [@corybrunson](#)

**Submitted:** 28 November 2019

**Published:** 07 January 2020

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC-BY](#)).

## Summary


Nanoparticles have attracted continued interest in academia and industry over the last few decades due to their remarkable properties that differ from the same materials in bulk. These properties are dependent on not only the size of the nanoparticles but also their shape. It is thus of great importance for nanoscientists to be able to predict the shape of nanoparticles of different materials and in different environments (Marks & Peng, 2016).

In the continuum (large particle) limit, the equilibrium shape of nanoparticles can be determined with the so-called Wulff construction (Wulff, 1901). Wulff's theorem states that the distance from the center of the nanoparticle to a facet is proportional to the surface energy of that facet. The equilibrium shape, often referred to as the Wulff shape, can thus be determined provided that the orientation-dependent surface energy is known.

The Wulff construction has been generalized to nanoparticles of icosahedral and decahedral geometry (Marks, 1983) as well as nanoparticles on surfaces (so-called Winterbottom constructions (Winterbottom, 1967)). The regular Wulff construction has been implemented in several software packages, including a submodule of the Python package `pymatgen` (Ong et al., 2013), a no longer maintained C++ package (Roosen, McCormack, & Carter, 1998), and a Wolfram Mathematica implementation with a graphical user interface (Zucker, Chatain, Dahmen, Hagège, & Carter, 2012). While the latter code has support for Winterbottom constructions, we have found no publicly available software that implements the icosahedral and decahedral Wulff construction. The aforementioned codes also seem to lack the ability to transform the created shapes into an atomistic representation, i.e., a nanoparticle of the Wulff shape consisting of atoms in a crystal structure, a feature of critical importance if the Wulff construction is to be used for atomistic simulations.

`WulffPack` is a Python package that carries out the Wulff construction and its generalizations using an efficient algorithm based on calculation of the convex hull (Barber, Dobkin, Dobkin, & Huhdanpaa, 1996) of the vertices of the dual of the Wulff polyhedron (Roosen et al., 1998; Virtanen et al., 2019). The user provides surface energies and crystal symmetry and `WulffPack` returns a versatile object that, at its core, contains the coordinates of the Wulff shape. Extraction of symmetry operations is handled internally with `spglib` (Togo & Tanaka, 2018). `WulffPack` includes functionality for visualizing the constructed shapes using `Matplotlib` (Hunter, 2007) (Fig. 1). There are also functions for analyzing the constructed shape, most notably in terms of area fraction of symmetrically inequivalent facets. This quantity is important in applications where properties of the material are facet-dependent, such as in catalysis. Finally, using the Atomic Simulation Environment (Larsen et al., 2017), an atomistic representation of the Wulff shape can also be extracted.

An extensive user guide including a documentation of the API is available at <http://wulffpack.materialsmodeling.org/>.



**Figure 1:** Truncated octahedron (left), decahedron (middle) and icosahedron (right), as visualized in WulffPack.

## Acknowledgements

This work was funded by the Knut and Alice Wallenberg Foundation and the Swedish Research Council.

## References

- Barber, C. B., Dobkin, D. P., Dobkin, D. P., & Huhdanpaa, H. (1996). The quickhull algorithm for convex hulls. *ACM Trans. Math. Softw.*, 22(4), 469–483. doi:[10.1145/235815.235821](https://doi.org/10.1145/235815.235821)
- Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. *Computing in Science & Engineering*, 9(3), 90–95. doi:[10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Larsen, A. H., Mortensen, J. J., Blomqvist, J., Castelli, I. E., Christensen, R., Duřak, M., Friis, J., et al. (2017). The Atomic Simulation Environment—A Python library for working with atoms. *Journal of Physics: Condensed Matter*, 29(27), 273002. doi:[10.1088/1361-648X/aa680e](https://doi.org/10.1088/1361-648X/aa680e)
- Marks, L. D. (1983). Modified Wulff constructions for twinned particles. *Journal of Crystal Growth*, 61(3), 556–566. doi:[10.1016/0022-0248\(83\)90184-7](https://doi.org/10.1016/0022-0248(83)90184-7)
- Marks, L. D., & Peng, L. (2016). Nanoparticle shape, thermodynamics and kinetics. *Journal of Physics: Condensed Matter*, 28(5), 053001. doi:[10.1088/0953-8984/28/5/053001](https://doi.org/10.1088/0953-8984/28/5/053001)
- Ong, S. P., Richards, W. D., Jain, A., Hautier, G., Kocher, M., Cholia, S., Gunter, D., et al. (2013). Python materials genomics (pymatgen): A robust, open-source python library for materials analysis. *Computational Materials Science*, 68, 314–319. doi:[10.1016/j.commatsci.2012.10.028](https://doi.org/10.1016/j.commatsci.2012.10.028)
- Roosen, A. R., McCormack, R. P., & Carter, W. C. (1998). Wulffman: A tool for the calculation and display of crystal shapes. *Computational Materials Science*, 11(1), 16–26. doi:[10.1016/S0927-0256\(97\)00167-5](https://doi.org/10.1016/S0927-0256(97)00167-5)
- Togo, A., & Tanaka, I. (2018). Spglib: A software library for crystal symmetry search. *arXiv e-prints*, arXiv:1808.01590. Retrieved from <http://arxiv.org/abs/1808.01590>
- Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., Burovski, E., et al. (2019). SciPy 1.0—Fundamental Algorithms for Scientific Computing in Python. *arXiv e-prints*, arXiv:1907.10121. Retrieved from <http://arxiv.org/abs/1907.10121>

- Winterbottom, W. L. (1967). Equilibrium shape of a small particle in contact with a foreign substrate. *Acta Metallurgica*, 15(2), 303–310. doi:[10.1016/0001-6160\(67\)90206-4](https://doi.org/10.1016/0001-6160(67)90206-4)
- Wulff, G. (1901). Zur Frage der Geschwindigkeit des Wachstums und der Auflösung der Krystallflächen. *Zeitschrift für Kristallographie*, 34, 449–530. doi:[10.1524/zkri.1901.34.1.449](https://doi.org/10.1524/zkri.1901.34.1.449)
- Zucker, R. V., Chatain, D., Dahmen, U., Hagège, S., & Carter, W. C. (2012). New software tools for the calculation and display of isolated and attached interfacial-energy minimizing particle shapes. *Journal of Materials Science*, 47(24), 8290–8302. doi:[10.1007/s10853-012-6739-x](https://doi.org/10.1007/s10853-012-6739-x)